

The Penrose inequality for an axially perturbed Schwarzschild black hole

J. Tafel

University of Warsaw

5-th PoToR conference
Wojanów 2018

joint work with J. Kopiński

The Penrose inequality

Area of the Schwarzschild horizon

$$A = 4\pi(2M)^2 = 16\pi M^2$$

hence

$$M = \sqrt{\frac{A}{16\pi}}.$$

For the Kerr horizon

$$A = 8\pi M(M + \sqrt{M^2 - a^2})$$

hence (the Penrose inequality)

$$M \geq \sqrt{\frac{A}{16\pi}}.$$

The Cauchy data with a horizon

If

- a future singularity is surrounded by the event horizon
(physically realistic data, the cosmic censorship conjecture)
- configuration tends to a stationary state

The Cauchy data with a horizon

If

- a future singularity is surrounded by the event horizon (physically realistic data, the cosmic censorship conjecture)
- configuration tends to a stationary state

then the no-hair theorem etc. (almost) imply that

- the end state is the Kerr metric with M_∞ and A_∞
- $M_\infty \leq M$ (because of radiation)
- $A_\infty \geq A$ (BH thermodynamics)

The Cauchy data with a horizon

If

- a future singularity is surrounded by the event horizon (physically realistic data, the cosmic censorship conjecture)
- configuration tends to a stationary state

then the no-hair theorem etc. (almost) imply that

- the end state is the Kerr metric with M_∞ and A_∞
- $M_\infty \leq M$ (because of radiation)
- $A_\infty \geq A$ (BH thermodynamics)

Conclusion: the Penrose inequality should be satisfied on the initial surface

Vacuum initial data with a horizon

Constraints on g_{ij}, K_{ij}

$$\nabla_i (K^{ij} - g^{ij} H) = 0$$

$$R + H^2 - K^2 = 0$$

where $H = K^i_i$ and $K^2 = K_{ij} K^{ij}$.

Vacuum initial data with a horizon

Constraints on g_{ij}, K_{ij}

$$\nabla_i (K^{ij} - g^{ij} H) = 0$$

$$R + H^2 - K^2 = 0$$

where $H = K^i_i$ and $K^2 = K_{ij}K^{ij}$.

Marginal outer trapped surface (MOTS) with unit normal vector n^i and mean curvature $h = \nabla_i n^i$:

$$\theta_+ = H - K_{nn} + h = 0$$

$$\theta_- = H - K_{nn} - h \leq 0.$$

Vacuum initial data with a horizon

Constraints on g_{ij}, K_{ij}

$$\nabla_i (K^{ij} - g^{ij} H) = 0$$

$$R + H^2 - K^2 = 0$$

where $H = K^i_i$ and $K^2 = K_{ij}K^{ij}$.

Marginal outer trapped surface (MOTS) with unit normal vector n^i and mean curvature $h = \nabla_i n^i$:

$$\theta_+ = H - K_{nn} + h = 0$$

$$\theta_- = H - K_{nn} - h \leq 0.$$

If $H = h = 0$ then the Penrose inequality follows from the Hamiltonian constraint (Geroch, ..., Huisken and Ilmanen).

The Lichnerowicz equation

Let $H = \nabla_i K^{ij} = 0$ and S_0 be a closed surface. Then

$$g'_{ij} = \psi^4 g_{ij}, \quad K'_{ij} = \psi^{-2} K_{ij}, \quad \psi > 0$$

satisfy the constraint equations if ψ satisfies the Lichnerowicz equation

$$\Delta \psi = \frac{1}{8} R \psi - \frac{1}{8} K^2 \psi^{-7}.$$

The Lichnerowicz equation

Let $H = \nabla_i K^{ij} = 0$ and S_0 be a closed surface. Then

$$g'_{ij} = \psi^4 g_{ij}, \quad K'_{ij} = \psi^{-2} K_{ij}, \quad \psi > 0$$

satisfy the constraint equations if ψ satisfies the Lichnerowicz equation

$$\Delta \psi = \frac{1}{8} R \psi - \frac{1}{8} K^2 \psi^{-7}.$$

If, in addition, ψ satisfies the boundary condition on S_0

$$n^i \partial_i \psi + \frac{1}{2} h \psi - \frac{1}{4} K_{nn} \psi^{-3} = 0$$

then S_0 becomes MOTS (D. Maxwell, extra conditions required).

Small perturbations of the Schwarzschild data

As the preliminary data we take

- the Schwarzschild initial data on $t = \text{const}$

$$g = \frac{dr^2}{1 - \frac{2M}{r}} + r^2(d\theta^2 + \sin^2 \theta d\varphi^2)$$

Small perturbations of the Schwarzschild data

As the preliminary data we take

- the Schwarzschild initial data on $t = \text{const}$

$$g = \frac{dr^2}{1 - \frac{2M}{r}} + r^2(d\theta^2 + \sin^2 \theta d\varphi^2)$$

- a nontrivial axially symmetric solution K of the momentum constraint with $H = 0$ (Conboye and O'Murchadha)

Small perturbations of the Schwarzschild data

As the preliminary data we take

- the Schwarzschild initial data on $t = \text{const}$

$$g = \frac{dr^2}{1 - \frac{2M}{r}} + r^2(d\theta^2 + \sin^2 \theta d\varphi^2)$$

- a nontrivial axially symmetric solution K of the momentum constraint with $H = 0$ (Conboye and O'Murchadha)
- the Schwarzschild horizon $r = 2M$ as the surface S_0

Small perturbations of the Schwarzschild data

As the preliminary data we take

- the Schwarzschild initial data on $t = \text{const}$

$$g = \frac{dr^2}{1 - \frac{2M}{r}} + r^2(d\theta^2 + \sin^2 \theta d\varphi^2)$$

- a nontrivial axially symmetric solution K of the momentum constraint with $H = 0$ (Conboye and O'Murchadha)
- the Schwarzschild horizon $r = 2M$ as the surface S_0

The Penrose inequality, if true, should follow from the Lichnerowicz equation and the boundary condition.

Approximation up to K^2 : $\psi = 1 + \psi_1 + \psi_2$

$$\Delta\psi_1 = 0, \quad n^i \partial_i \psi_1 = \frac{1}{4} K_{nn} \text{ on } S_0 \quad (1)$$

Approximation up to K^2 : $\psi = 1 + \psi_1 + \psi_2$

$$\Delta\psi_1 = 0, \quad n^i \partial_i \psi_1 = \frac{1}{4} K_{nn} \text{ on } S_0 \quad (1)$$

$$\Delta\psi_2 = -\frac{1}{8} K_{ij} K^{ij}, \quad n^i \partial_i \psi_2 = -\frac{3}{4} K_{nn} \psi_1 \text{ on } S_0 \quad (2)$$

Approximation up to K^2 : $\psi = 1 + \psi_1 + \psi_2$

$$\Delta\psi_1 = 0, \quad n^i \partial_i \psi_1 = \frac{1}{4} K_{nn} \text{ on } S_0 \quad (1)$$

$$\Delta\psi_2 = -\frac{1}{8} K_{ij} K^{ij}, \quad n^i \partial_i \psi_2 = -\frac{3}{4} K_{nn} \psi_1 \text{ on } S_0 \quad (2)$$

Total energy (after the conformal transformation):

$$E = M - \frac{1}{2\pi} \int_{S_\infty} \partial_r \psi d\sigma.$$

Area of the horizon:

$$A = \int_{S_0} \psi^4 d\sigma.$$

An integration of the Lichnerowicz equation yields

$$E = M + \frac{1}{16\pi} \int_{S_0}^{\infty} K^2 dV - \frac{1}{8\pi} \oint_{S_0} K_{nn}(1 - 3\psi_1) d\sigma$$

and a more subtle integration yields a similar expression for A .

An integration of the Lichnerowicz equation yields

$$E = M + \frac{1}{16\pi} \int_{S_0}^{\infty} K^2 dV - \frac{1}{8\pi} \oint_{S_0} K_{nn}(1 - 3\psi_1) d\sigma$$

and a more subtle integration yields a similar expression for A .

The Penrose inequality is equivalent to

$$\int_{2M}^{\infty} r^2 \langle K^2 \rangle dr + 8M^3 \langle K_{nn} \rangle^2 - 12M \langle \psi_1^2 \rangle \geq 0 \quad (3)$$

where $\langle \dots \rangle$ denotes integral over the unit sphere and ψ_1 depends on K_{nn} via the Neumann problem.

An integration of the Lichnerowicz equation yields

$$E = M + \frac{1}{16\pi} \int_{S_0}^\infty K^2 dV - \frac{1}{8\pi} \oint_{S_0} K_{nn}(1 - 3\psi_1) d\sigma$$

and a more subtle integration yields a similar expression for A .

The Penrose inequality is equivalent to

$$\int_{2M}^\infty r^2 \langle K^2 \rangle dr + 8M^3 \langle K_{nn} \rangle^2 - 12M \langle \psi_1^2 \rangle \geq 0 \quad (3)$$

where $\langle \dots \rangle$ denotes integral over the unit sphere and ψ_1 depends on K_{nn} via the Neumann problem.

Problem

Given $K_{nn} \geq 0$ on S_0 what is minimal value of the first integral in (3).

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .
 Q' does not appear in K_{nn} and generates terms enlarging K^2 .

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .
 Q' does not appear in K_{nn} and generates terms enlarging K^2 .
Dependence on Q

$$K_{r\theta} = -\frac{Q_{,r}}{r \sin \theta}, \quad K_{rr} = \frac{Q_{,\theta}}{r^3 \sin \theta}$$

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .
 Q' does not appear in K_{nn} and generates terms enlarging K^2 .
Dependence on Q

$$K_{r\theta} = -\frac{Q_{,r}}{r \sin \theta}, \quad K_{rr} = \frac{Q_{,\theta}}{r^3 \sin \theta}$$

$$(K_{\theta\theta} \sin^2 \theta)_{,\theta} = (rQ_{,r})_{,r} \sin \theta - \frac{\cos \theta}{r} Q_{,\theta}$$

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .
 Q' does not appear in K_{nn} and generates terms enlarging K^2 .
Dependence on Q

$$K_{r\theta} = -\frac{Q_{,r}}{r \sin \theta}, \quad K_{rr} = \frac{Q_{,\theta}}{r^3 \sin \theta}$$

$$(K_{\theta\theta} \sin^2 \theta)_{,\theta} = (rQ_{,r})_{,r} \sin \theta - \frac{\cos \theta}{r} Q_{,\theta}$$

$$K_{\varphi}^{\varphi} = -K_r^r - K_{\theta}^{\theta}.$$

Parametrization of axially symmetric solutions K_{ij}

Such solutions of $\nabla_i K^{ij} = 0$ depend on 2 functions Q, Q' of r and θ .
 Q' does not appear in K_{nn} and generates terms enlarging K^2 .

Dependence on Q

$$K_{r\theta} = -\frac{Q_{,r}}{r \sin \theta}, \quad K_{rr} = \frac{Q_{,\theta}}{r^3 \sin \theta}$$

$$(K_{\theta\theta} \sin^2 \theta)_{,\theta} = (rQ_{,r})_{,r} \sin \theta - \frac{\cos \theta}{r} Q_{,\theta}$$

$$K_{\varphi}^{\varphi} = -K_r^r - K_{\theta}^{\theta}.$$

A convenient form of Q :

$$Q = \sin \theta q_{,\theta} - a \cos \theta, \quad a = \text{const}$$

where q is regular and $q_{,r}$ is bounded at ∞ .

Expansion into Legendre polynomials $P_n(z)$, $z = \cos \theta$:

$$q = \sum_{n=1} q_n(r) P_n.$$

$K_{\theta\theta}$ creates problems in $\langle K^2 \rangle$.

Expansion into Legendre polynomials $P_n(z)$, $z = \cos \theta$:

$$q = \sum_{n=1} q_n(r) P_n.$$

$K_{\theta\theta}$ creates problems in $\langle K^2 \rangle$.

Miracle No 1

$$K_{\theta\theta} + \frac{1}{2}r^2 K_{rr} = \sum_{n=2} ((rq_{n,r})_{,r} - \frac{1}{2}n(n+1)\frac{q_n}{r}) \tilde{P}_n,$$

where polynomials

$$\tilde{P}_n = \frac{1}{n+2} (nP_n - \frac{2}{n-1} P_{n-1,z})$$

are orthogonal.

Expansion into Legendre polynomials $P_n(z)$, $z = \cos \theta$:

$$q = \sum_{n=1} q_n(r) P_n.$$

$K_{\theta\theta}$ creates problems in $\langle K^2 \rangle$.

Miracle No 1

$$K_{\theta\theta} + \frac{1}{2} r^2 K_{rr} = \sum_{n=2} ((r q_{n,r})_{,r} - \frac{1}{2} n(n+1) \frac{q_n}{r}) \tilde{P}_n,$$

where polynomials

$$\tilde{P}_n = \frac{1}{n+2} (n P_n - \frac{2}{n-1} P_{n-1,z})$$

are orthogonal.

Hence $\langle K^2 \rangle = \sum F_n(q_n)$, $F_n \geq 0$ and we can look separately for a minimum of each integral

$$\int_{2M}^{\infty} r^2 F_n dr.$$

Miracle No 2

Variation of integrals leads to fourth order linear ordinary equations for q_n , $n \geq 2$ which can be exactly solved (with a help of Mathematica).

Miracle No 2

Variation of integrals leads to fourth order linear ordinary equations for q_n , $n \geq 2$ which can be exactly solved (with a help of Mathematica).

Solutions read $q_n = a_n + \frac{b_n}{r^n} + \frac{c_n}{r^{n+1}}$ where constants are defined by P_n components of K_{rr} on the horizon.

Miracle No 2

Variation of integrals leads to fourth order linear ordinary equations for q_n , $n \geq 2$ which can be exactly solved (with a help of Mathematica).

Solutions read $q_n = a_n + \frac{b_n}{r^n} + \frac{c_n}{r^{n+1}}$ where constants are defined by P_n components of K_{rr} on the horizon.

Now minimal value of the integrals can be calculated. The Penrose inequality follows except the case $q = (br + \frac{c}{r}) \cos \theta$ (then the 4-th order approximation in K yields the inequality).

Miracle No 2

Variation of integrals leads to fourth order linear ordinary equations for q_n , $n \geq 2$ which can be exactly solved (with a help of Mathematica).

Solutions read $q_n = a_n + \frac{b_n}{r^n} + \frac{c_n}{r^{n+1}}$ where constants are defined by P_n components of K_{rr} on the horizon.

Now minimal value of the integrals can be calculated. The Penrose inequality follows except the case $q = (br + \frac{c}{r}) \cos \theta$ (then the 4-th order approximation in K yields the inequality).

Main result

The Penrose inequality is preserved under a small addition of axially symmetric exterior curvature to the Schwarzschild data.

Miracle No 2

Variation of integrals leads to fourth order linear ordinary equations for q_n , $n \geq 2$ which can be exactly solved (with a help of Mathematica).

Solutions read $q_n = a_n + \frac{b_n}{r^n} + \frac{c_n}{r^{n+1}}$ where constants are defined by P_n components of K_{rr} on the horizon.

Now minimal value of the integrals can be calculated. The Penrose inequality follows except the case $q = (br + \frac{c}{r}) \cos \theta$ (then the 4-th order approximation in K yields the inequality).

Main result

The Penrose inequality is preserved under a small addition of axially symmetric exterior curvature to the Schwarzschild data.

First (to my knowledge) proof with horizon which is not a minimal surface.

Generalizations: nonsymmetric K , nonspherical horizons (?)